

Logic (PHIL 2080, COMP 2620, COMP 6262)
Chapter: Propositional Logic — Semantic Tableaux

Pascal Bercher

AI Group
School of Computing
College of Engineering and Computer Science
the Australian National University

“14 March” 2022 (Canberra day! Pre-recording)



Australian
National
University

Introduction

Recap: This course so far!

- What's a formula:
 - Syntactically (i.e., how do they look like)
 - Semantically (i.e., what do they *mean*, which properties do they have; keywords: *Interpretations*, *Satisfiability*, *Unsatisfiability*, *Tautology*)

Recap: This course so far!

- What's a formula:
 - Syntactically (i.e., how do they look like)
 - Semantically (i.e., what do they *mean*, which properties do they have; keywords: *Interpretations*, *Satisfiability*, *Unsatisfiability*, *Tautology*)
- What's a sequent:
 - Syntactically (i.e., $X \vdash A$ and $X \Vdash A$)
 - Semantically (i.e., $X \vdash A$ is called *valid*, $X \Vdash A$, if each interpretation that makes all formulae in X true also makes A true.)

Recap: This course so far!

- What's a formula:
 - Syntactically (i.e., how do they look like)
 - Semantically (i.e., what do they *mean*, which properties do they have; keywords: *Interpretations*, *Satisfiability*, *Unsatisfiability*, *Tautology*)
- What's a sequent:
 - Syntactically (i.e., $X \vdash A$ and $X \Vdash A$)
 - Semantically (i.e., $X \vdash A$ is called *valid*, $X \Vdash A$, if each interpretation that makes all formulae in X true also makes A true.)
- How to prove validity?
 - With truth tables (that's the definition; but takes too long)
 - With Natural deduction (often much quicker, but 'harder')

Today: Motivation

- When you were asked to prove $X \vdash A$ with Natural Deduction (ND), then... you were able to do so! The proof existed!
- Why? Because you were only proving *valid* sequents!
- Why is that problematic?

Today: Motivation

- When you were asked to prove $X \vdash A$ with Natural Deduction (ND), then... you were able to do so! The proof existed!
- Why? Because you were only proving *valid* sequents!
- Why is that problematic?
 - Because you cannot *decide* validity with ND!
 - Suppose somebody asks: Is $X \vdash A$ valid, what do you do?
 - You can attempt ND, but if you fail: then why? Did you just not try hard enough? Or isn't it possible?

Today: Motivation

- When you were asked to prove $X \vdash A$ with Natural Deduction (ND), then... you were able to do so! The proof existed!
 - Why? Because you were only proving *valid* sequents!
 - Why is that problematic?
 - Because you cannot *decide* validity with ND!
 - Suppose somebody asks: Is $X \vdash A$ valid, what do you do?
 - You can attempt ND, but if you fail: then why? Did you just not try hard enough? Or isn't it possible?
 - Today: We learn a second proof system, which cannot only prove validity (if it's valid), but it can also *disprove* validity (if it's invalid)!
 - We call this: *deciding validity*.
- This is the Semantic Tableaux proof system!

Recap on Definitions

- Hopefully everyone recalls the meaning of $X \models A$:
It means that A logically follows from the formulae in X , i.e., that sequent is *valid*, which is defined in terms of truth tables:
 - Each interpretation that makes all formulae in X true also also makes A true.
 - Or: There is no interpretation that makes X true, but not A .

Recap on Definitions

- Hopefully everyone recalls the meaning of $X \models A$:
It means that A logically follows from the formulae in X , i.e., that sequent is *valid*, which is defined in terms of truth tables:
 - Each interpretation that makes all formulae in X true also also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- So what did $X \vdash A$ mean again?
 - It was actually just *short* for $X \vdash_{ND} A$.
 - It meant: A can be derived from X using Natural Deduction.
 - Yoshi will show that this implies validity, but technically it was just referring to Natural Deduction and manipulating formulae.

Recap on Definitions

- Hopefully everyone recalls the meaning of $X \models A$:
It means that A logically follows from the formulae in X , i.e., that sequent is *valid*, which is defined in terms of truth tables:
 - Each interpretation that makes all formulae in X true also also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- So what did $X \vdash A$ mean again?
 - It was actually just *short* for $X \vdash_{ND} A$.
 - It meant: A can be derived from X using Natural Deduction.
 - Yoshi will show that this implies validity, but technically it was just referring to Natural Deduction and manipulating formulae.
- Today, we learn how to decide validity using Semantic Tableaux.
 - Depending on context, $X \vdash A$ might stand for either Natural Deduction ($X \vdash_{ND} A$) or Semantic Tableaux ($X \vdash_{ST} A$).

Semantic Tableaux Proof Idea

General Idea behind Semantic Tableaux

- Semantic tableaux has its name because its proof technique mirrors/directly exploits the definition of validity of a sequent.
- So recall what $X \models A$ means:
 - Each interpretation that makes all formulae in X true also makes A true.
 - Or: There is no interpretation that makes X true, but not A .

General Idea behind Semantic Tableaux

- Semantic tableaux has its name because its proof technique mirrors/directly exploits the definition of validity of a sequent.
- So recall what $X \models A$ means:
 - Each interpretation that makes all formulae in X true also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- We pursue *proof by contradiction* to exploit this definition!
- General idea: Assume the sequent is invalid and detect a contradiction. From this contradiction we can infer that our assumption of invalidity must be wrong, and we can conclude validity.
- An additional advantage: If we don't get a contradiction we can even prove invalidity! (Which Natural Deduction can't!)

Semantic Tableaux as “Proof by Contradiction”

- For the fourth time today (sorry...), what's validity?
 - Each interpretation that makes all formulae in X true also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- So what's that “inverted” property, i.e., *not valid* or *invalid*?

Semantic Tableaux as “Proof by Contradiction”

- For the fourth time today (sorry...), what's validity?
 - Each interpretation that makes all formulae in X true also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- So what's that “inverted” property, i.e., *not* valid or *invalid*?
 - There exists an interpretation such that:
 - ▶ it makes all formulae in X true
 - ▶ but it does *not* make A true!

Semantic Tableaux as “Proof by Contradiction”

- For the fourth time today (sorry...), what's validity?
 - Each interpretation that makes all formulae in X true also makes A true.
 - Or: There is no interpretation that makes X true, but not A .
- So what's that “inverted” property, i.e., *not* valid or *invalid*?
 - There exists an interpretation such that:
 - ▶ it makes all formulae in X true
 - ▶ but it does *not* make A true!
 - If that leads to a contradiction, the sequent was valid!
 - If we *find* such an interpretation, we have a counterexample! :)

Examples for the Proof Idea (Example 1, which happens to be valid)

- Suppose we want to decide the sequent $\overbrace{p \wedge q}^X \vdash \overbrace{q \vee \neg p}^A$
- So we assume invalidity, meaning:
 - We assume there's an interpretation that makes $p \wedge q$ true
 - and the same interpretation does not make $q \vee \neg p$ true.

p	q	$\overbrace{p \wedge q}^X$	$\overbrace{q \vee \neg p}^A$
0	0	0	1
0	1	0	1
1	0	0	0
1	1	1	1

Examples for the Proof Idea (Example 1, which happens to be valid)

- Suppose we want to decide the sequent $\overbrace{p \wedge q}^X \vdash \overbrace{q \vee \neg p}^A$
- So we assume invalidity, meaning:
 - We assume there's an interpretation that makes $p \wedge q$ true
 - and the same interpretation does not make $q \vee \neg p$ true.

p	q	$\overbrace{p \wedge q}^X$	$\overbrace{q \vee \neg p}^A$
0	0	0	1
0	1	0	1
1	0	0	0
1	1	1	1

There's only one interpretation that makes $p \wedge q$ true, namely $I(p) = 1$ and $I(q) = 1$, but this one *also* makes $q \vee \neg p$ true!

- So our assumption that such an interpretation exists was wrong!
- So such an interpretation does not exist.
- But that's the definition of validity! :) So it's valid!

Examples for the Proof Idea (Example 2, which (also) happens to be valid)

- Suppose we want to decide the sequent $\overbrace{p \vee q}^X \vdash \overbrace{p \vee \neg p}^A$
- So we assume invalidity, meaning:
 - We assume there's an interpretation that makes $p \vee q$ true
 - and the same interpretation does not make $p \vee \neg p$ true.

p	q	$\overbrace{p \vee q}^X$	$\overbrace{p \vee \neg p}^A$
0	0	0	1
0	1	1	1
1	0	1	1
1	1	1	1

There are many interpretations that make $p \vee q$ true, but all of them also make $p \vee \neg p$ true!

- So our assumption that such an interpretation exists was wrong!
- So such an interpretation does not exist.
- But that's (still) the definition of validity, so it's valid!

Examples for the Proof Idea (Example 3, which happens to be *invalid*)

- Suppose we want to decide the sequent $\overbrace{p \vee q}^X \vdash \overbrace{p \wedge q}^A$
- So we assume invalidity, meaning:
 - We assume there's an interpretation that makes $p \vee q$ true
 - and the same interpretation does not make $p \wedge q$ true.

p	q	$\overbrace{p \vee q}^X$	$\overbrace{p \wedge q}^A$
0	0	0	0
0	1	1	0
1	0	1	0
1	1	1	1

Examples for the Proof Idea (Example 3, which happens to be *invalid*)

- Suppose we want to decide the sequent $\overbrace{p \vee q}^X \vdash \overbrace{p \wedge q}^A$
- So we assume invalidity, meaning:
 - We assume there's an interpretation that makes $p \vee q$ true
 - and the same interpretation does not make $p \wedge q$ true.

p	q	$\overbrace{p \vee q}^X$	$\overbrace{p \wedge q}^A$
0	0	0	0
0	1	1	0
1	0	1	0
1	1	1	1

Yes, now we found two cases!

$I_1(p) = 0, I_1(q) = 1$ makes $p \vee q$ true, but not $p \wedge q$. Same for $I_2(p) = 1, I_2(q) = 0$!

- So we found *two* interpretations that *prove* invalidity! (One is sufficient, though.)
- Well... That's it! We proved invalidity.

Proof Idea, Concluding Comments

- Bad news: That was just the *proof idea*!
- The actual *proof technique* does *not* use truth tables.
- After all: proof techniques were developed so that we do not need them! Recall that proof tables scale exponentially with the number of propositional variables!
- These examples were just to illustrate:
 - How that proof by contradiction idea works here
 - Why and how Semantic Tableaux acts as decision procedure

The Proof Technique in more Detail

- Maintain a list of formulae, label each either **T** (*true*) or **F** (*false*).
- Initially, label all formulae in X with **T**, and A with **F**.

The Proof Technique in more Detail

- Maintain a list of formulae, label each either **T** (*true*) or **F** (*false*).
- Initially, label all formulae in X with **T**, and A with **F**.
- Then simplify each formula and flip truth values as required. E.g.,
 - If some line holds **F** : $q \vee \neg p$, we get two more: **F** : q and **F** : $\neg p$
 - If some line holds **F** : $\neg p$, we get another line: **T** : p

The Proof Technique in more Detail

- Maintain a list of formulae, label each either **T** (*true*) or **F** (*false*).
- Initially, label all formulae in X with **T**, and A with **F**.
- Then simplify each formula and flip truth values as required. E.g.,
 - If some line holds **F** : $q \vee \neg p$, we get two more: **F** : q and **F** : $\neg p$
 - If some line holds **F** : $\neg p$, we get another line: **T** : p
- All lines below each other “belong together” and define one shared interpretation. Some rules will *branch*, i.e., create another set of lines existing “in parallel”. So different branches describe different interpretations.
- Once no more formulae can be simplified (and hence all propositional symbols have a truth value assigned), we either:
 - 1 Have obtained a consistent (contradiction-free) interpretation that proves invalidity. (Such a branch is called **open branch**.)
 - 2 Or if each branch leads to a contradiction (e.g., **T** : p and **F** : p – or even with some formula), we proved validity.

How to Support Climate Change – or: How to Prove (In)Validity

$$p \vee q, \neg p \vdash q$$

$$\begin{array}{l} (1) \quad \mathbf{T}: \quad p \vee q \\ (2) \quad \mathbf{T}: \quad \neg p \end{array} \quad \left. \vphantom{\begin{array}{l} (1) \\ (2) \end{array}} \right\} \text{premises } X$$

How to Support Climate Change – or: How to Prove (In)Validity

$$p \vee q, \neg p \vdash q$$

- | | | | | |
|-----|-----------|------------|---|----------------|
| (1) | T: | $p \vee q$ | } | premises X |
| (2) | T: | $\neg p$ | | |
| (3) | F: | q | } | conclusion A |

- Within each path, all lines “accumulate”.

How to Support Climate Change – or: How to Prove (In)Validity

$$p \vee q, \neg p \vdash q$$

- (1) **T:** $p \vee q$
- (2) **T:** $\neg p$ ✓
- (3) **F:** q
- (4) **F:** p from (2)

- Within each path, all lines “accumulate”.

How to Support Climate Change – or: How to Prove (In)Validity

$$p \vee q, \neg p \vdash q$$

(1) **T:** $p \vee q$ ✓

(2) **T:** $\neg p$ ✓

(3) **F:** q

(4) **F:** p from (2)

(5) **T:** p ⚡ from (1)

(6) **T:** q ⚡ from (1)

- Within each path, all lines “accumulate”.
- Branches branch, i.e., they split different possibilities.
- If all leafs die, the tree dies: Success! *Sequent is valid.*

How to Support Climate Change – or: How to Prove (In)Validity

$$p \vee q, \neg p \vdash q$$

$$(1) \quad \mathbf{T}: \quad p \vee q \quad \checkmark$$

$$(2) \quad \mathbf{T}: \quad \neg p \quad \checkmark$$

$$(3) \quad \mathbf{F}: \quad q$$

$$(4) \quad \mathbf{F}: \quad p \quad \text{from (2)}$$

$$(5) \quad \mathbf{T}: \quad p \quad \text{from (1)}$$

$$(6) \quad \mathbf{T}: \quad q \quad \text{from (1)}$$

- Within each path, all lines “accumulate”.
- Branches branch, i.e., they split different possibilities.
- If all leaves die, the tree dies: Success! *Sequent is valid*.
- If some leaf survives, the tree lives: Failure!
 - The sequent is invalid.
 - We can extract an interpretation invalidating the sequent.

All Simplification Rules

Rule Set (And, Or, Not)

And Elimination:

$$\underline{\mathbf{T}: A \wedge B}$$

$$\underline{\mathbf{F}: A \wedge B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Rule Set (And, Or, Not)

And Elimination:

$$\frac{\mathbf{T}: A \wedge B}{\mathbf{T}: A, \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \wedge B}{\mathbf{F}: A, \mathbf{F}: B}$$

The comma means that both lines hold in the same branch, i.e., we can write them below each other!

<i>A</i>	<i>B</i>	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Rule Set (And, Or, Not)

And Elimination:

$$\frac{\mathbf{T}: A \wedge B}{\mathbf{T}: A, \mathbf{T}: B}$$

$$\mathbf{T}: A, \mathbf{T}: B$$

$$\frac{\mathbf{F}: A \wedge B}{\mathbf{F}: A \mid \mathbf{F}: B}$$

$$\mathbf{F}: A \mid \mathbf{F}: B$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

The comma means that both lines hold in the same branch, i.e., we can write them below each other!

The bar (“|”) means that we branch over different possibilities, so the lines end up in different branches!

Rule Set (And, Or, Not)

And Elimination:

$$\frac{\mathbf{T}: A \wedge B}{\mathbf{T}: A, \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \wedge B}{\mathbf{F}: A \mid \mathbf{F}: B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Or Elimination:

$$\frac{\mathbf{T}: A \vee B}{\quad}$$

$$\frac{\mathbf{F}: A \vee B}{\quad}$$

A	B	\vee
0	0	0
0	1	1
1	0	1
1	1	1

Rule Set (And, Or, Not)

And Elimination:

$$\frac{\mathbf{T}: A \wedge B}{\mathbf{T}: A, \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \wedge B}{\mathbf{F}: A \mid \mathbf{F}: B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Or Elimination:

$$\frac{\mathbf{T}: A \vee B}{\mathbf{T}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \vee B}{\quad}$$

A	B	\vee
0	0	0
0	1	1
1	0	1
1	1	1

Rule Set (And, Or, Not)

And Elimination:

$$\frac{T: A \wedge B}{T: A, T: B}$$

$$\frac{F: A \wedge B}{F: A \mid F: B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Or Elimination:

$$\frac{T: A \vee B}{T: A \mid T: B}$$

$$\frac{F: A \vee B}{F: A, F: B}$$

A	B	\vee
0	0	0
0	1	1
1	0	1
1	1	1

Rule Set (And, Or, Not)

And Elimination:

$$\frac{T: A \wedge B}{T: A, T: B}$$

$$\frac{F: A \wedge B}{F: A \mid F: B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Or Elimination:

$$\frac{T: A \vee B}{T: A \mid T: B}$$

$$\frac{F: A \vee B}{F: A, F: B}$$

A	B	\vee
0	0	0
0	1	1
1	0	1
1	1	1

Negation Elimination:

$$\frac{T: \neg A}{\quad}$$

$$\frac{F: \neg A}{\quad}$$

A	\neg
0	1
1	0

Rule Set (And, Or, Not)

And Elimination:

$$\frac{\text{T: } A \wedge B}{\text{T: } A, \text{ T: } B}$$

$$\frac{\text{F: } A \wedge B}{\text{F: } A \mid \text{F: } B}$$

A	B	\wedge
0	0	0
0	1	0
1	0	0
1	1	1

Or Elimination:

$$\frac{\text{T: } A \vee B}{\text{T: } A \mid \text{T: } B}$$

$$\frac{\text{F: } A \vee B}{\text{F: } A, \text{ F: } B}$$

A	B	\vee
0	0	0
0	1	1
1	0	1
1	1	1

Negation Elimination:

$$\frac{\text{T: } \neg A}{\text{F: } A}$$

$$\frac{\text{F: } \neg A}{\text{T: } A}$$

A	\neg
0	1
1	0

Rule Set (Implication)

Implication Elimination:

$$\frac{\mathbf{T}: A \rightarrow B}{}$$
$$\frac{\mathbf{F}: A \rightarrow B}{}$$

A	B	\rightarrow
0	0	1
0	1	1
1	0	0
1	1	1

Rule Set (Implication)

Implication Elimination:

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{}$$

A	B	\rightarrow
0	0	1
0	1	1
1	0	0
1	1	1

Rule Set (Implication)

Implication Elimination:

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

A	B	\rightarrow
0	0	1
0	1	1
1	0	0
1	1	1

Rule Set (Implication)

Implication Elimination:

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

A	B	\rightarrow
0	0	1
0	1	1
1	0	0
1	1	1

- Note that $X \vdash A$ iff $X \models A$ intuitively holds, because these rules mimic the truth tables *exactly*.

Rule Set (Implication)

Implication Elimination:

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

A	B	\rightarrow
0	0	1
0	1	1
1	0	0
1	1	1

- Note that $X \vdash A$ iff $X \models A$ intuitively holds, because these rules mimic the truth tables *exactly*.
- Also keep in mind that we only write $X \vdash A$ instead of $X \vdash_{ND} A$ or $X \vdash_{ST} A$ since the applied proof system is clear from context.

Examples!

Example (for a valid Sequent)

$$\vdash p \rightarrow (q \rightarrow p)$$

$$(1) \quad \mathbf{F}: p \rightarrow (q \rightarrow p)$$

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

Example (for a valid Sequent)

$$\vdash p \rightarrow (q \rightarrow p)$$

- (1) **F:** $p \rightarrow (q \rightarrow p)$ ✓
(2) **T:** p from (1)
(3) **F:** $q \rightarrow p$ from (1)

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

Example (for a valid Sequent)

$$\vdash p \rightarrow (q \rightarrow p)$$

- (1) **F:** $p \rightarrow (q \rightarrow p)$ ✓
- (2) **T:** p from (1)
- (3) **F:** $q \rightarrow p$ ✓ from (1)
- (4) **T:** q from (3)
- (5) **F:** p ⚡ from (3)

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

- This sequent is valid, because all branches show a contradiction!

Example (for a valid Sequent)

$$\vdash p \rightarrow (q \rightarrow p)$$

- (1) **F:** $p \rightarrow (q \rightarrow p)$ ✓
- (2) **T:** p from (1)
- (3) **F:** $q \rightarrow p$ ✓ from (1)
- (4) **T:** q from (3)
- (5) **F:** p ⚡ from (3)

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

- This sequent is valid, because all branches show a contradiction!
- Here, there was no branching. Normally, we have ≥ 2 branches, and *all* have to show a contradiction.
- We also did not use any assumptions here (that would have been labeled *true* (**T**)), because there weren't any.

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ (1) **T:** $p \rightarrow (q \vee r)$ (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

$\mathbf{T}: A \rightarrow B$
$\mathbf{F}: A \mid \mathbf{T}: B$

$\mathbf{F}: A \rightarrow B$
$\mathbf{T}: A, \mathbf{F}: B$

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ from (2)

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

$$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$$

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ from (2)
 (5) **F:** $s \rightarrow \neg q$ from (3)

$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$
--

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$
--

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ from (3)
 (6) **T:** $p \wedge s$ from (4)
 (7) **F:** r from (4)

$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$
--

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$
--

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ ✓ from (3)
 (6) **T:** $p \wedge s$ from (4)
 (7) **F:** r from (4)
 (8) **T:** s from (5)
 (9) **F:** $\neg q$ from (5)

$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$
--

$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$
--

$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$
--

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ ✓ from (3)
 (6) **T:** $p \wedge s$ ✓ from (4)
 (7) **F:** r from (4)
 (8) **T:** s from (5)
 (9) **F:** $\neg q$ from (5)
 (10) **T:** p from (6)

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

$$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$$

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ ✓ from (3)
 (6) **T:** $p \wedge s$ ✓ from (4)
 (7) **F:** r from (4)
 (8) **T:** s from (5)
 (9) **F:** $\neg q$ ✓ from (5)
 (10) **T:** p from (6)
 (11) **T:** q from (9)

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

$$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$$

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$ ✓
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ ✓ from (3)
 (6) **T:** $p \wedge s$ ✓ from (4)
 (7) **F:** r from (4)
 (8) **T:** s from (5)
 (9) **F:** $\neg q$ ✓ from (5)
 (10) **T:** p from (6)
 (11) **T:** q from (9)

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

$$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$$

(12) **F:** p ⚡ from (1)

(13) **T:** $q \vee r$ from (1)

Example (for an Invalid Sequent)

Attempting to show validity of $p \rightarrow (q \vee r) \vdash \neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$

- (1) **T:** $p \rightarrow (q \vee r)$ ✓
 (2) **F:** $\neg(s \rightarrow \neg q) \rightarrow ((p \wedge s) \rightarrow r)$ ✓
 (3) **T:** $\neg(s \rightarrow \neg q)$ ✓ from (2)
 (4) **F:** $(p \wedge s) \rightarrow r$ ✓ from (2)
 (5) **F:** $s \rightarrow \neg q$ ✓ from (3)
 (6) **T:** $p \wedge s$ ✓ from (4)
 (7) **F:** r from (4)
 (8) **T:** s from (5)
 (9) **F:** $\neg q$ ✓ from (5)
 (10) **T:** p from (6)
 (11) **T:** q from (9)

$$\frac{\mathbf{T}: A \rightarrow B}{\mathbf{F}: A \mid \mathbf{T}: B}$$

$$\frac{\mathbf{F}: A \rightarrow B}{\mathbf{T}: A, \mathbf{F}: B}$$

$$\frac{\mathbf{T}: \neg A}{\mathbf{F}: A}$$

Interpretation: $I(r) = 0, I(s) = I(q) = I(p) = 1$

- (12) **F:** p \downarrow from (1)

- (13) **T:** $q \vee r$ ✓ from (1) $\left\langle \begin{array}{l} (14) \mathbf{T}: q \text{ open! from (13)} \\ (15) \mathbf{T}: r \downarrow \text{ from (13)} \end{array} \right.$

Comments About Previous Proof(s)

- In the beginning (when only the first lines were shown) we had the choice of which implication to simplify.
- We chose line (2), because its rule does not branch, and it's always good to postpone branching as long as possible so we don't "duplicate" work.

Comments About Previous Proof(s)

- In the beginning (when only the first lines were shown) we had the choice of which implication to simplify.
- We chose line (2), because its rule does not branch, and it's always good to postpone branching as long as possible so we don't "duplicate" work.
- We detected an *open branch*, i.e., a complete path where no further reductions were possible.

Comments About Previous Proof(s)

- In the beginning (when only the first lines were shown) we had the choice of which implication to simplify.
- We chose line (2), because its rule does not branch, and it's always good to postpone branching as long as possible so we don't "duplicate" work.
- We detected an *open branch*, i.e., a complete path where no further reductions were possible.
 - Via "collecting" the truth assignments to atoms along the open branch we can construct an interpretation.

Comments About Previous Proof(s)

- In the beginning (when only the first lines were shown) we had the choice of which implication to simplify.
- We chose line (2), because its rule does not branch, and it's always good to postpone branching as long as possible so we don't "duplicate" work.
- We detected an *open branch*, i.e., a complete path where no further reductions were possible.
 - Via "collecting" the truth assignments to atoms along the open branch we can construct an interpretation.
 - That interpretation is a witness that:
 - ▶ that there exists an interpretation that makes X true but not A ,
 - ▶ and, thus (by definition), that A does not follow logically from X .
 - ▶ This means that the sequent is *invalid*.

Comments About Previous Proof(s)

- In the beginning (when only the first lines were shown) we had the choice of which implication to simplify.
- We chose line (2), because its rule does not branch, and it's always good to postpone branching as long as possible so we don't "duplicate" work.
- We detected an *open branch*, i.e., a complete path where no further reductions were possible.
 - Via "collecting" the truth assignments to atoms along the open branch we can construct an interpretation.
 - That interpretation is a witness that:
 - ▶ that there exists an interpretation that makes X true but not A ,
 - ▶ and, thus (by definition), that A does not follow logically from X .
 - ▶ This means that the sequent is *invalid*.
- Also note that we have a contradiction whenever some *formula* appears true and false within the same branch. We do not need to wait until it is atomic.

Summary

Content of this Lecture

- We covered *Semantic Tableaux*, which “mimics” the definition of validity.
 - All rules required to simplify formulae as required.
 - You learned (or realized) that:
 - Natural Deduction cannot *decide* validity.
 - Semantic Tableaux *can* decide validity.
- We covered the entire Logic Notes sections:
- More about propositional logic: Semantic tableaux